

## Casirivimab (a.k.a. REGN10933) Injection (120 mg/ml)

### MUST ADMINISTER WITH IMDEVIMAB (a.k.a. REGN10987)

**Manufacturer:** Regeneron Pharmaceuticals, Inc.

**Distributor:** AmerisourceBergen (a.k.a. ABC);  
[c19therapies@amerisourcebergen.com](mailto:c19therapies@amerisourcebergen.com)


**Product:** Component of 2-drug monoclonal antibody cocktail for outpatient infusion

**Emergency Use Authorization (EUA):** 11/21/20

**Units:** Supplied in 2 volumes

- 1332 mg/11.1 ml single-dose vial in carton (10 ml from 1 vial needed for patient course)
- 300 mg/2.5 ml single-dose vials in cartons (10 ml; 2.5 ml from each of 4 vials needed for patient course)

*Note: Some cartons and vials of casirivimab may be instead labeled REGN10933*



**Example:** 1332 mg/11.1 ml carton



**Example:** 300 mg/2.5 ml carton

**Storage:** Keep in carton until use. Unopened vials must be stored at refrigerated temperature (2°C–8°C / 36°F–46°F) until use. Do not freeze, shake, or expose to direct light.

**Earliest Expiration Date of Units Shipped:** 06/2022

**Single Vial w/ Carton Dimensions:** 1.7”(d) x 1.8”(w) x 2.8”(h)

**Single Vial w/ Carton Weight:** 1.5 oz. for 1332 mg/11.1 ml vial; 0.9 oz. for 300 mg/2.5 ml vial

**Case:** A full case contains 24 vials in cartons

**Case Dimensions:** 9.6”(d) x 7.4”(w) x 4.6”(h)

**Case Weight:** 2.7 lb. for 1332mg/11.1 ml vials; 1.8 lb. for 300 mg/2.5 ml vials

**Administration:** (For [detailed guidelines](#), click here)

1. Inject 10 ml (1200 mg) casirivimab
2. Inject 10 ml (1200 mg) imdevimab
3. Infuse final volume (270 ml) containing cocktail of 2 monoclonal antibodies over at least 60 min

*Note: If immediate patient infusion is not possible, store the diluted cocktail solution at refrigerated temperature (2°C–8°C / 36°F–46°F) for no more than 36 hours and at room temperature for no more than 4 hours, including infusion time.*

**Resources/Links:**

- [Regeneron webpage](#), including:
  - [FDA Letter of Authorization](#)
  - [Fact Sheet for Healthcare Providers](#)
  - [Fact Sheet for Patients and Caregivers \(English\)](#)
  - [Fact Sheet for Patients and Caregivers \(Spanish\)](#)
- [FDA Frequently Asked Questions on the EUA for Casirivimab + Imdevimab](#)
  - [FAQ: Allocation, Distribution, and Administration of Casirivimab + Imdevimab](#)

## Imdevimab (a.k.a. REGN10987) Injection (120 mg/ml) MUST ADMINISTER WITH CASIRIVIMAB (a.k.a. REGN10933)

**Manufacturer:** Regeneron Pharmaceuticals, Inc.

**Distributor:** AmerisourceBergen (a.k.a. ABC);  
[c19therapies@amerisourcebergen.com](mailto:c19therapies@amerisourcebergen.com)


**Product:** Component of 2-drug monoclonal antibody cocktail for outpatient infusion

**Emergency Use Authorization (EUA):** 11/21/20

**Units:** Supplied in 2 volumes

- 1332 mg/11.1 ml single-dose vial in carton (10 ml from 1 vial needed for patient course)
- 300 mg/2.5 ml single-dose vials in cartons (10 ml; 2.5 ml each from 4 vials needed for patient course)

*Note: Some cartons and vials of imdevimab may be instead labeled REGN10987*



**Example:** 1332 mg/11.1 ml carton



**Example:** 300 mg/2.5 ml carton

**Storage:** Keep in carton until use. Unopened vials must be stored at refrigerated temperature (2°C–8°C / 36°F–46°F) until use. Do not freeze, shake, or expose to direct light.

**Earliest Expiration Date of Units Shipped:** 06/2022

**Single Vial w/ Carton Dimensions:** 1.7”(d) x 1.8”(w) x 2.8”(h)

**Single Vial w/ Carton Weight:** 1.5 oz. for 1332 mg/11.1 ml vial; 0.9 oz. for 300 mg/2.5 ml vial

**Case:** A full case contains 24 vials in cartons

**Case Dimensions:** 9.6”(d) x 7.4”(w) x 4.6”(h)

**Case Weight:** 2.7 lb. for 1332mg/11.1 ml vials; 1.8 lb. for 300 mg/2.5 ml vials

**Administration:** (For [detailed guidelines](#), click here)

1. Inject 10 ml (1200 mg) casirivimab
2. Inject 10 ml (1200 mg) imdevimab
3. Infuse final volume (270 ml) containing cocktail of 2 monoclonal antibodies over at least 60 min

*Note: If immediate patient infusion is not possible, store the diluted cocktail solution at refrigerated temperature (2°C–8°C / 36°F–46°F) for no more than 36 hours and at room temperature for no more than 4 hours, including infusion time.*

**Resources/Links:**

- [Regeneron webpage](#), including:
  - [FDA Letter of Authorization](#)
  - [Fact Sheet for Healthcare Providers](#)
  - [Fact Sheet for Patients and Caregivers \(English\)](#)
  - [Fact Sheet for Patients and Caregivers \(Spanish\)](#)
- [FDA Frequently Asked Questions on the EUA for Casirivimab + Imdevimab](#)
- [FAQ: Allocation, Distribution, and Administration of Casirivimab + Imdevimab](#)