

Table 5: How Casirivimab and Imdevimab are Supplied

Antibody	Concentration	Package Size	NDC Number
Casirivimab REGN 10933	1332 mg/11.1 mL (120 mg/mL)	1 vial per pack	61755-024-01
	300 mg/2.5 mL (120 mg/mL)	1 vial per pack	61755-026-01
Imdevimab REGN10987	1332 mg/11.1 mL (120 mg/mL)	1 vial per pack	61755-025-01
	300 mg/2.5 mL (120 mg/mL)	1 vial per pack	61755-027-01

Storage and Handling

Casirivimab is preservative-free. Discard any unused portion.

Imdevimab is preservative-free. Discard any unused portion.

Store unopened casirivimab and imdevimab vials in a refrigerator at 2°C to 8°C (36°F to 46°F) in the original carton to protect from light.

DO NOT FREEZE. DO NOT SHAKE. DO NOT EXPOSE TO DIRECT LIGHT.

Solution in vial requires dilution prior to administration. The prepared infusion solution is intended to be used immediately. If immediate administration is not possible, store diluted casirivimab and imdevimab solution in the refrigerator at 2°C to 8°C (36°F to 46°F) for no more than 36 hours and at room temperature up to 25°C (77°F) for no more than 4 hours, including infusion time. If refrigerated, allow the infusion solution to equilibrate to room temperature for approximately 30 minutes prior to administration.